

Our thanks are due to Professor T. H. Johnston, who kindly allowed the section cutting to be performed in the laboratory of the Biology Dept., University of Queensland.

#### BIBLIOGRAPHY.

- BAILEY, F. M., Queensland Flora, 1901, Pt. IV., pp. 1351-1352.  
 BENTHAM, G., Flora Australiensis, 1870, V., p. 532.  
 LONGMAN AND WHITE, "Mutation in a Proteaceous Tree." Proc. Roy. Soc. Q'land., XXX., 1918, pp. 162-4.  
 MUELLER, F. v. Fragm. Phytog., 1868, VI., pp. 247-8.

#### EXPLANATION OF TEXT-FIGURES.

Transverse sections across flowers of *Buckinghamia celsissima* F.v.M.

Fig. 1.—Mutant of Longman and White; *c*, deeply stained cells in parenchyma of stipes and aborted styles; *h1*, *h2*, *h3*, segments of hypogynous gland; *p*, perianth segments; *par*, parenchyma in stipes; *s*, stipes; *st. 1*, *st. 2*, aborted styles; *v.b.*, vascular bundles.

Fig. 2.—Normal flower; lettering similar to above; *h*, undivided, semi-annular hypogynous gland.

---

#### ALTERATION OF GENERIC NAME.

---

NOTE BY J. DOUGLAS OGILBY.

---

In these Proceedings, Vol. xxi, p. 91, I proposed the name *Eurycaulus* for a genus of belonoid fishes. This, having been previously used in Coleoptera by Fairmaire, 1868, I now change to *Tropidocaulus*.